

This product contains no phosphorous compounds. Effective yet economical, this product cleans, disinfects and deodorizes in one labor saving step. This product is a one-step disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, the AIDS Virus and Hepatitis B Virus and Hepatitis C Virus), fungicidal, and inhibits the growth of mold and mildew, and their odors, when used as directed.

This product is effective at eliminating odors on porous surfaces such as upholstery, drapes, carpets, bedding, shower curtains, mattresses and non-porous surfaces such as walls and floors. Excellent for fire restoration. Use to remove odors caused by flooding and smoke from fires.

Use in: airplanes, airports, transportation terminals, boats, buses, cars, emergency vehicles, ships, taxis, trailers, trains, food storage areas, public rest rooms, dressing rooms, exercise facilities, locker rooms, shower and bath areas, institutional and public facilities.

Use on washable hard, nonporous surfaces of: bathroom fixtures, cabinets, coils and drain pans of air conditioning and refrigeration equipment and heat pumps, chairs, countertops, desks, floors, garbage cans, exteriors of refrigerators, refrigerated storage and display equipment, showers, sinks, stovetops, tables, toilets, urinals, walls, other hard nonporous surfaces made of: glazed ceramic, glazed enameled surfaces, glazed porcelain, laminated surfaces, metal, plastic (such as polystyrene or polypropylene), sealed stone, stainless steel, vinyl and plastic upholstery and finished woodwork.

This product is a one-step disinfectant, Bactericidal according to the current AOAC Use-Dilution Test Method and Virucidal* according to the virucidal qualification modified in the presence of 200 ppm hard water plus 5% organic serum against: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica*, *Escherichia coli*, *Salmonella typhi*, *Staphylococcus aureus*-Methicillin-Resistant (MRSA), *Staphylococcus aureus*-Multi-Drug Resistant, *Staphylococcus aureus*-Vancomycin Intermediate Resistant (VISA), *Streptococcus pyogenes*, *Hepatitis B Virus (HBV), *Hepatitis C Virus (HCV), *Herpes Simplex Virus Type 1, *Herpes Simplex Virus Type 2, *HIV-1 (AIDS Virus), *Human Coronavirus, *Influenza Virus Type A/Brazil, *Norwalk Virus- Feline Calicivirus, *Respiratory Syncytial Virus (RSV), *SARS Associated Coronavirus, *Vaccinia Virus, *Aspergillus niger*, *Trichophyton mentagrophytes*.

This product is a sanitizer for hard nonporous non-food contact surfaces according to the method for Sanitizers - Non-Food Contact Surfaces, modified in the presence of 500 ppm hard water plus 5% organic serum in 60 seconds against:

Klebsiella pneumoniae *Staphylococcus aureus*

Dilution:

Disinfection (1:256)½ oz. per gallon of water.....848 ppm active
 Disinfection (1:128)1 oz. per gallon of water.....1695 ppm active
 Disinfection (1:64)2 oz. per gallon of water.....3390 ppm active
 Disinfection (1:16)8 oz. per gallon of water.....13562 ppm active
 Sanitizer (1:1024)½ oz. per 4 gallons of water...200 ppm active quat

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

If you do not understand the label, find someone to explain it to you in detail.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

Preparation of Use-Solution: Mix ½ oz. per gallon of water (1:256). For heavy duty use, mix 2 oz. per gallon of water (1:64).

DISINFECTION/VIRUCIDAL*/FUNGICIDAL/MOLD AND MILDEW CONTROL DIRECTIONS:

Apply use-solution to hard, nonporous surfaces, thoroughly wetting surfaces with a cloth, mop, sponge, sprayer or by immersion. Treated surfaces must remain wet for 10 minutes.

For Influenza Virus Type A and Human Coronavirus, treated surfaces must remain wet for 1 minute. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from the surface; rub with a brush, sponge or cloth. Do not breathe spray. Rinse all surfaces that come in contact with food such as countertops, exteriors of appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

For fungicidal activity, this product is an effective fungicide against *Trichophyton mentagrophytes* [the Athlete's Foot Fungus] in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. Dilute at ½ oz. per gallon in 200 ppm hard water and allow surface to remain wet for 10 minutes.

For mold and mildew: At ½ oz. per gallon, this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

Prepare a fresh solution daily or more often if the use solution becomes visibly diluted or soiled.

This product is not for use on medical device surfaces.

TO CLEAN AND DEODORIZE: Mix ½ oz. of this product per gallon of water to clean and deodorize surfaces. Apply using a cloth, mop, sponge or sprayer. Wipe or allow to air dry.

Deodorizing directions: Nursing Homes, Hospitals, Hotels, Schools, Restaurants: Using ½ - 2 oz. in chemical toilets, waste receptacles, bed pans, drainage bottles, diaper pails, hampers, disposals, commodes and air conditioner pans will eliminate odors.

Mopping Solutions, Automatic Scrubbers: Using ½ - 2 oz. per gallon of cleaning solution eliminates undesirable odors.

Carpets: ½ - 1 oz. or ½ - 2 oz. per gallon of shampoo solution eliminates odors associated with urine, vomit, smoke, and mildew. Reapply after cleaning for freshening effects. Allow to dry.

Non-Acid Toilet Bowl and Urinal Disinfectant / Cleaner Directions: Remove gross filth prior to disinfection.

From Concentrate: Add ½ oz. to the bowl and mix. Brush over exposed surfaces and under the rim, allow to stand for 10 minutes and flush.

From Use Solution: Empty toilet bowl or urinal and apply ½ oz. per gallon use-solution to exposed surfaces and under the rim with a cloth, mop, sponge or mechanical spray. Brush or swab thoroughly; allow to stand for 10 minutes and flush.

For Heavy Duty Use: Empty toilet bowl or urinal and apply 2 oz. per gallon use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly, allow to stand for 10 minutes and flush.

FOR USE ON FINISHED FLOORS: To limit gloss reduction use this product at ½ oz. per gallon. Apply with a damp mop or autocrusher. Allow surface to remain wet for 10 minutes. Wipe or allow to air dry. Do not use on floor finish at 2 oz. per gallon.

***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of Zep Triton.

Contact time: Leave surface wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV with ½ oz. per gallon use-solution. Use a 10-minute contact time for disinfection against all other viruses, bacteria and fungi claimed.

Disposal of infectious material: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

TO SANITIZE NON-FOOD CONTACT SURFACES: Mix ½ oz. of this product per 4 gallons of water to sanitize hard nonporous surfaces. Treated surfaces must remain wet for 60 seconds. Wipe dry with a sponge, mop, or cloth or allow to air dry. Do not use in food preparation, food handling and food storage areas. This product is not to be used on glassware, dishware, cookware or utensils.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and reclose the container. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat two more times. Then offer for recycling if available

or puncture and dispose in sanitary landfill or by other procedures approved by state and local authorities. Follow pesticide disposal instructions for rinsate. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations.

11/18

EPA REG. NO. 6836-78-1270
 EPA EST. NO. 1270-TX-1(2) or 40849-GA-1(E)
 Postscript is first letter or number of batch code on container.



MANUFACTURED IN USA
A Zep Inc. Brand
 Distributed by:
 Zep Inc.
 350 Joe Frank Harris Parkway, SE
 Emerson, GA 30137
 www.zep.com
 1-877-1-BUY-ZEP/1-877-428-9937

V202-3



TRITON

One-Step Disinfectant • Sanitizer • Fungicide • Mildewstat • Virucide* • Deodorizer for Hospitals, Institutional and Industrial Use

• Staphylocidal • Pseudomonacidal • Bactericidal • Salmonellacidal • Fungicidal • Virucidal*

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride	6.510%
Diocetyl dimethyl ammonium chloride.....	2.604%
Didecyl dimethyl ammonium chloride	3.906%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	8.680%
OTHER INGREDIENTS:	78.300%
TOTAL	100.000%

**KEEP OUT OF REACH OF CHILDREN
 DANGER • PELIGRO**

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective (rubber or chemical resistant) gloves. Harmful if swallowed and/or if absorbed through the skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

Net Contents: 1 GALLON (3.785 liters)